

## Common Medicine Dosages

**\*\*Do Not use Ibuprofen in babies less than 6 months unless instructed by physician\*\***

**1 tsp = 5 ml    \*\*A cc and ml are the same thing\*\***

**Benadryl dosages are for allergic reaction. Use half dosage for runny nose**

<i>Childs Weight</i>		<i>Benadryl</i>	<i>Acetomniophen</i>	<i>Ibuprofen</i>	<i>Mucinex Multi</i>
<i>Kg</i>	<i>Pounds</i>	<i>12.5 mg/ ml</i>	<i>160 mg/5ml</i>	<i>100 mg/5ml</i>	<i>Symptom Cold</i>
1	2.2	0.4	0.5		
2	4.4	0.8	1		
3	6.6	1.2	1.5		
4	8.8	1.6	2		
5	11	2	2.5	2.5	
6	13.2	2.4	3	3	1.2
7	15.4	2.8	3.5	3.5	1.4
8	17.6	3.2	4	4	1.6
9	19.8	3.6	4.5	4.5	1.8
10	22	4	5	5	2
11	24.2	4.4	5.5	5.5	2.2
12	26.4	4.8	6	6	2.4
13	28.6	5.2	6.5	6.5	2.6
14	30.8	5.6	7	7	2.8
15	33	6	7.5	7.5	3
16	35.2	6.4	8	8	3.2
17	37.4	6.8	8.5	8.5	3.4
18	39.6	7.2	9	9	3.6
19	41.8	7.6	9.5	9.5	3.8
20	44	8	10	10	4
21	46.2	8.4	10.5	10.5	4.2
22	48.4	8.8	11	11	4.4
23	50.6	9.2	11.5	11.5	4.6
24	52.8	9.6	12	12	4.8
25	55	10	12.5	12.5	5
26	57.2	10.4	13	13	5.2
27	59.4	10.8	13.5	13.5	5.4
28	61.6	11.2	14	14	5.6
29	63.8	11.6	14.5	14.5	5.8
30	66	12	15	15	6
31	68.2	12.4	15.5	15.5	6.2
32	70.4	12.8	16	16	6.4
33	72.6	13.2	16.5	16.5	6.6
34	74.88	13.6	17	17	6.8
35	77	14	17.5	17.5	7
36	79.2	14.4	18	18	7.2
37	81.4	14.8	18.5	18.5	7.4
38	83.6	15.2	19	19	7.6
39	85.8	15.6	19.5	19.5	7.8
40	88	16	20	20	8